



**University
Hospitals Sussex**
NHS Foundation Trust

Coping with diarrhoea

Patient information

Diarrhoea involves the passing of frequent, loose stools. It may be caused by an infection or by treatment which is irritating your bowel.

Diarrhoea can also be accompanied by loss of appetite, nausea, vomiting, stomach pains, high temperature and dehydration.

Short-term modification of your diet can sometimes help to relieve symptoms.

If your diarrhoea is severe or persists, consult your doctor.

Fluid

It is very important to maintain a high fluid intake to prevent dehydration.

- Aim for at least 1.5 to 2 litres (3 pints) of fluid per day
- Clear fluids are likely to be better tolerated, e.g. fruit squashes, cordials, tea, decaffeinated coffee, Bovril, clear soups or Oxo

Ready mixed rehydration sachets can be purchased from chemists and added to drinking water. These are useful for replacing lost fluid and salt. Follow the manufacturers' recommendations.

Try the following

- Small frequent meals and snacks
- Chew food well and eat slowly
- Relax after meals.
- Choose soft, easily digested foods, e.g. white fish, mashed potatoes, sorbets, jelly, yoghurt, milk puddings, ice creams
- Choose low fibre varieties of food, e.g. white bread, white pasta and rice, plain biscuits and low fibre cereals such as Cornflakes or Rice Krispies.

Avoid the following

These foods may aggravate your diarrhoea or cause abdominal cramps or excess wind/gas:

- High fibre foods, e.g. wholemeal bread, wholegrain cereals, brown rice and pasta and foods made with wholemeal flour
- Fruit juice
- Nuts and dried fruits
- Uncooked vegetables and fruit
- Beans, lentils and pulses
- Cauliflower, cabbage and onions
- Spicy, fried or very fatty foods, e.g. curries, chillies, cream, pastry, chips, etc.
- Very hot or very cold food and drinks
- Alcohol, strong tea or coffee
- Avoid any food which you know upsets you
- Fatty food.

Remember

If your diarrhoea is severe or persists, consult your doctor.

Contact us

For further information or to provide feedback

St Richard's Hospital

Telephone: **01243 831498**

Email: uhsussex.chichesterdietitians@nhs.net

Worthing & Southlands Hospital

Telephone: **01903 286779**

Email: worthing.dietitians@nhs.net

St Richard's Hospital
Spitalfield Lane
Chichester, West Sussex
PO19 6SE

Southlands Hospital
Upper Shoreham Road
Shoreham-by-Sea
West Sussex
BN43 6QT

We are committed to making our publications as accessible as possible. If you need this document in an alternative format, for example, large print, Braille or a language other than English, please contact the Communications Office by emailing uhsussex.communications@nhs.net

www.uhsussex.nhs.uk

Reference no. 2195
Department: Dietetics
Issue date: 03/23
Review date: 03/26
Author: University Hospital Sussex Bariatric Dietitians
Version: 3

© University Hospitals Sussex NHS Foundation Trust Disclaimer: The information in this leaflet is for guidance purposes only and is in no way intended to replace professional clinical advice by a qualified practitioner.

